Elective Lymph Node Dissection in Malignant Melanoma

Dr. Janix M. De Guzman Presentor

Premise

In patients with malignant melanoma with a clinically negative lymph node,

Is lymph node dissection indicated?

Lymph node dissection

Goals:

- accurate staging?
- improve survival rate?
- prevent spread?

The Benefits

- Lymph node involvement is defined in AJCC as:
 - N1 metastasis 3 cm or less in greatest dimension in any regional lymph node
 - N2 metastasis more than 3 cm in greatest dimension in any regional lymph node and/or in-transit metastasis

AJCC staging System for Melanoma, Showing Reported 5-Year Survival Rates for Each Stage¹

AJCC stage	TNM categories	5-year survival rate (%)
	T1N0M0	
	T2N0M0	90
	T3N0M0	
II	T4N0M0	70
III	Any T, Any N, M0	50
IV	Any T, Any N, M1	20

5-year survival rate of clinically negative but pathologically positive nodes ²⁻³

50%

prospective randomized trials (level 1A evidences) failed to show a benefit for elective lymph node dissection in terms of survival 4-6

	# of patients	Study groups	stratification	Significance demonstrated In survival rate	% change identified in trial
Veronesi, et al	553	ELND Delayed	none	none	none
Cascinelli, et al	252	ELND observe	Gender, site histology	none	none
Balch, et al	740	ELND observe	Age histology	none	none

Breslow, measured depth of invasion – predictor of nodal metastasis⁷

Breslow's thickness	Likelihood of nodal metastasis	5-year survival (clinically negative node)
In situ	0	100%
<u><</u> 0.75 mm	5%	96%
0.76 – 1.5 mm	10-20%	87%
1.51 – 4 mm	25-30%	70%
>4 mm	70%	47%

What can we offer after adequate staging?

	# of patients	Study groups	stratificati on	Significance demonstrated
Creagan ⁸	232	IFN observe	none	none
Grob ⁹	499	IFN control	none	Relapse-free interval
Pehambe rger ¹⁰	311	IFN 4 observe	none	Disease free interval
Kirkwood 11	642	High dose Low dose observation	Stage nodes	none

Regarding Spread

- ELND may not prevent recurrent nodal disease in the dissected basin.¹²
- Melanoma spreads via both lymphatic channels and the blood stream.
- Some patients will develop distant metastasis without manifesting clinical node involvement. For deeper (>4mm) lesions 70% chance of distant metastasis.¹³

The Risks¹²

- Seroma
- Infection
- Skin Necrosis
- Lymphedema
 - Once it develops, no effective cure. Treatment are available that palliate symptoms, but these must be continued for the patients lifetime.

Morbidity

 Difficult to justify in patient who has undergone lymph node dissection with no histologically positive nodes.

Elective lymph node dissection, Is it indicated? NO.

Reasons in Summary:

- absence of any apparent impact on survival among patients who underwent ELND
- 2. no available effective adjuvant treatment
- 3. May not prevent recurrent nodal disease in the dissected basin.
- 4. Patients subjected to significant risks without concrete benefit.

References

- 1. Chang AE, Karnell LH, Menck HR. the National Cancer Data Base report on cutaneous and non-cutaneous melanoma: a summary of 84,836 cases from the past decade. Cancer 1998;83:1664-1678
- 2. Reintgen D, Balch CM, Kirkwood J, Ross M. Recent advances in the care of the patient with malignant melanoma. Ann Surg 1997;225:1.
- 3. Harris MN, shapiro RL, Roses DF. Malignant Melanoma, primary surgical management (excision and node dissection) based on pathology and staging. Cancer 1995;75:715.

- 4. Veronesi U, Adamus J, Bandiera DC, et al. Delayed regional lymph node dissection in stage I melanoma of the skin of the lower extremities. Cancer. 1982;49(11):2420-30.
- 5. Cascinelli N, Morabito A, Santinami M, MacKie RM, Belli F. Immediate or delayed dissection of regional nodes in patients with melanoma of the trunk: a randomised trial. WHO Melanoma Programme. Lancet. 1998;351(9105):793-6.
- 6. Balch CM, Soong SJ, Smith T, et al. Investigators from the Intergroup Melanoma Surgical Trial. Long-term results of a prospective surgical trial comparing 2 cm vs. 4 cm excision margins for 740 patients with 1-4 mm melanomas. Ann Surg Oncol. 2001;8(2):101-8.

- 7. Breslow A. Thickness, cross-sectional area and depth of invasion in the prognosisof cutaneous melanoma. Ann Surg. 1970;172:982
- Creagan ET, Dalton RJ, Ahmann DL, et al. Randomized, surgical adjuvant clinical trial of recombinant interferon alfa-2a in selected patients with malignant melanoma. J Clin Oncol. 1995;13(11):2776-83.
- 9. Grob JJ, Dreno B, de la Salmoniere P, et al. Randomised trial of interferon alpha-2a as adjuvant therapy in resected primary melanoma thicker than 1.5 mm without clinically detectable node metastases. French Cooperative Group on Melanoma. Lancet. 1998, 27;351(9120):1905-10.

- 10. Pehamberger H, Soyer HP, Steiner A, et al. Adjuvant interferon alfa-2a treatment in resected primary stage II cutaneous melanoma. Austrian Malignant Melanoma Cooperative Group. J Clin Oncol. 1998;16(4):1425-9.
- 11. Kirkwood JM, Strawderman MH, Ernstoff MS, Smith TJ, Borden EC, Blum RH. Interferon alfa-2b adjuvant therapy of high-risk resected cutaneous melanoma: the Eastern Cooperative Oncology Group Trial EST 1684. J Clin Oncol. 1996;14(1):7-17.
- 12. Slingluff CL Jr, Stidham KR, Ricci WM, Stanley WE, Seigler HF. Surgical management of regional lymph nodes in patients with melanoma. Experience with 4682 patients. Ann Surg. 1994;219(2):120-30.

- 13. Reintgen DS, Cox EB, McCarty KS, Vollmer RT, Seigler HF. Efficacy of elective lymph node dissection in patients with intermediate thickness primary melanoma. Ann Surg. 1983;198:339
- 14. Strobbe LJ, Jonk A, Hart AA, Nieweg OE, Kroon BB. Positive iliac and obturator nodes in melanoma: survival and prognostic factors. Ann Surg Oncol. 1999;6(3):255-62